


**Tools for Practice** is proudly sponsored by the Alberta College of Family Physicians (ACFP). ACFP is a provincial, professional voluntary organization, representing more than 3,800 family physicians, family medicine residents and medical students in Alberta. Established over fifty years ago, the ACFP strives for excellence in family practice through advocacy, continuing medical education and primary care research. [www.acfp.ca](http://www.acfp.ca)

November 18, 2013



## **NSAIDs and cardiovascular safety: the truth makes my heart hurt**

**Clinical Question: Do different non-steroidal anti-inflammatory drugs (NSAIDs) have different cardiovascular risks?**

**Bottom-line: COX-2 inhibitors and traditional NSAIDs, except naproxen, increase the risk of major vascular events and death. When prescribing NSAIDs, patients' gastrointestinal (GI) and cardiovascular (CV) risk should be assessed, with naproxen or low-dose ibuprofen preferentially chosen for patients at risk of CV disease.**

### **Evidence:**

- Meta-analysis of 754 randomized controlled trials (~350,000 patients):<sup>1</sup>
  - Mixed population, primarily arthritis, at low-moderate CV risk (i.e. CV event rate ~1% per year).
  - COX-2 inhibitors, compared to placebo, increased:
    - All-cause mortality, Rate Ratio (RR) 1.22 (95% Confidence Interval 1.04–1.44).
    - Major CV events, RR 1.37 (1.14–1.66).
  - Diclofenac (150 mg/day): similar risks to COX-2s for mortality [RR 1.02 (0.84–1.24)] and CV events [RR 0.97 (0.84–1.12)].
    - Indirectly, diclofenac significantly increases CV events [RR 1.41 (1.12–1.78)] but not mortality [RR 1.20 (0.94–1.54)] compared to placebo.
  - Naproxen (1000 mg/day) has less CV events and mortality than COX-2 inhibitors and may be similar to placebo.
  - Relative risk similar between patients with or without prior CV disease.
- Meta-analysis of observational trials:<sup>2</sup>
  - All COX-2s and NSAIDs, except naproxen and low-dose ibuprofen ( $\leq 1200$  mg/day), increase CV risk.
  - Risk increases with increasing NSAID dose.
- Results are consistent with previous meta-analysis.<sup>3</sup>

### **Context:**

- In Canada, naproxen (28%), celecoxib (21%), and diclofenac (17%) account for the majority of NSAIDs prescribed.<sup>4</sup>

- The magnitude of the CV risk with high-risk NSAIDs is similar to the magnitude of the CV benefit with statin therapy. Choosing high-risk NSAIDs (taken daily) can cause one additional CV event over five years in:<sup>1</sup>
  - ~100 low-risk patients (baseline 5% ten-year CV risk).
  - ~25 high-risk patients (baseline 20% ten-year CV risk).
- Generally, NSAIDs with relatively lower CV risks (naproxen) have relatively higher GI complication risks (ulcers and bleeds) and vice versa (coxibs).<sup>5</sup>
  - Adding a proton pump inhibitor to a non-selective NSAID results in similar GI complication risks as COX-2s.<sup>6</sup>
- All NSAIDs increase risk of heart failure.<sup>1</sup>

#### Authors:

Megan Harbin BScPharm ACPR, Ricky D Turgeon BScPharm ACPR,  
Michael R Kolber BSc MD CCFP MSc

#### References:

1. Coxib and traditional NSAID Trialists' (CNT) Collaboration. *Lancet*. 2013; 382(9894): 769–79.
2. McGettigan P, Henry D. *PLoS Med*. 2011; 8(9):1–18.
3. Trelle S, Reichenbach S, Wandel S, *et al*. *BMJ*. 2011; 342:c7086.
4. McGettigan P, Henry D. *PLoS Med*. 2013; 10(2):1–6.
5. Castellsague J, Riera-Guardia N, Calingaert, *et al*. *Drug Saf*. 2012; 35(12):1127–46.
6. Wang X, Tian HJ, Yang HK, *et al*. *Eur J Gastroenterol Hepatol*. 2011; 23:876–80.

**Tools for Practice** is a biweekly article summarizing medical evidence with a focus on topical issues and practice modifying information. It is coordinated by G. Michael Allan, MD, CCFP and the content is written by practising family physicians who are joined occasionally by a health professional from another medical specialty or health discipline. Each article is peer-reviewed, ensuring it maintains a high standard of quality, accuracy, and academic integrity. If you are not a member of the ACFP and would like to receive the TFP emails, please sign up for the distribution list at <http://bit.ly/signupfortfp>. Archived articles are available on the ACFP website.

This communication reflects the opinion of the authors and does not necessarily mirror the perspective and policy of the Alberta College of Family Physicians.